

D-instantons probing D3-branes and the AdS/CFT correspondence.

E.T. Akhmedov¹

Institute of Theoretical and Experimental Physics
Moscow, 117259, B. Cheremushkinskaya, 25.

Abstract

D-instantons are considered as a probe of coinciding N D3-branes. They can feel an external metric via the commutator terms in their effective action. We show that when the D-instantons are separated from the D3-branes, the metric which is probed at the one loop level, *exactly* coincides with that of the BPS R-R 3-brane. Interesting connection of this result to the possible explanation of the AdS/CFT correspondence within IKKT M-atrrix theory is discussed.

Dedicated to the memory of Alexander Valerievich Makedonski.

1. One of the ways of thinking about super-string theory is that it suggests a method of probing the geometry of an ambient space. Until recently we have had only one object which was able to serve for that purpose. It was the string itself. At present, however, at our disposal there are D-branes [1] which also can play the role of probes.

Remarkably, the D-branes can feel a geometry at distances smaller than the string scale [2]. This way it is possible to probe the curved space in the vicinity of string solitons [3, 4]. In some circumstances that is relevant for the investigation of the SYM low energy dynamics [5, 3]. For example, the D-branes give a geometric description of different phenomena in the SYM theories which live on their world-volumes (see, for example, [3, 5, 6, 7]).

Besides that, among various discoveries within the context of the D-brane physics, there is the AdS/CFT correspondence [8]. It establishes a relation between two seemingly different theories. One of them is the four-dimensional $\mathcal{N} = 4$, $SU(N)$ SYM theory, taken at large N and strong coupling ($g_{YM}^2 N = g_s N \gg 1$). While the other is the type IIB SUGRA on the $AdS_5 \times (S_5)_N$ background. The latter contains N units of the elementary flux of the R-R tensor field through the S_5 , which is indicated by the subscript " N ". Also

¹e-mail: akhmedov@vxitep.itep.ru

the radii of the AdS_5 and S_5 are equivalent to each other and given by $R^4 = 4\pi g_s N \alpha'^2$. Here g_s and α' are the string coupling constant and the inverse string tension, respectively.

Inspired by the developments of the probe physics we want to study how it can help in the explanation of the AdS/CFT correspondence. For this purpose we solve a warmup exercise which we hope could clear up the physics underlying the correspondence. Following [9], where a D-string was studied as a probe of an M5-brane, we consider n D-instantons probing N coinciding D3-branes. Via these tools the theory describing the M5-brane was examined within M-atrrix model [9]. Here we are doing the same for the D3-brane within the IKKT picture of the M-atrrix theory [10, 11].

The D-instantons in the background of the D3-branes have several branches on their "vacuum moduli space"². Only two of them are relevant for our discussion. In accordance with the ref. [9], the first one we refer to as "Coulomb branch". It corresponds to the case when all the D-instantons are separated from the coinciding D3-branes. The second one is referred to as "Higgs branch" [9]. It, in turn, describes the situation when all the D-instantons are sitting on the D3-branes. Of particular interest for our purposes are the effective instanton actions for the "massless fields" on the branches.

Concretely, in this note we calculate the one loop effective D-instanton action on the "Coulomb branch". As in the ref. [4], we want to find the geometry which is probed in this context. Below we explain that the D-instanton can feel the metric of a bulk space via the *commutator terms* in its effective action.

At first sight, our calculation is valid at fixed N and weak coupling, then ³ $g_{YM}^2 N = g_s N \ll 1$. However, because of the aid of SUSY we exactly recover the D3-brane metric, on the branch in question. This is the main result of our note.

Moreover, because the dilaton of the 3-brane SUGRA solution is just a constant, in our case it is possible to understand the singular limit of a small distance between the probe and the D3-brane. In this limit we get the theory describing the vicinity of the point on the D-instanton "vacuum moduli space", which connects the "Coulomb" and "Higgs" branches. Inspired by the ref. [9], we believe that its examination is relevant for the explanation of the AdS/CFT correspondence. At least within IKKT M-atrrix model.

In fact, first, the theory on the "Coulomb branch" at the singular point can be obtained when $N \rightarrow \infty$ and $g_s N \gg 1$ [9, 8]. Accordingly, *from our calculation* in this limit we get the D-instanton theory on the $AdS_5 \times S_5$ background.

Second, in accordance with the ref. [9], it is the latter theory which should be equivalent to that on the "Higgs branch". In our case this can be traced as follows. Really, the n D-instantons on the "Higgs branch" are equivalent to the ordinary n YM instantons of the $U(N)$ gauge theory [13, 14]. At the same time it is known that the moduli space of one $\mathcal{N} = 4$ super-instanton is exactly $AdS_5 \times S_5$ [15, 16, 17, 18]. Moreover, as was shown in [17], when $N \rightarrow \infty$, the n instanton moduli space is again the $AdS_5 \times S_5$. (All instantons tend to sit at the same point!) Hence, the latter should be the manifold which is seen by the D-instanton on the "Higgs branch".

²It is worth mentioning at this point that in the case of D-instantons there is no such a thing as the vacuum moduli space. Rather the manifold in question corresponds to the minimization of the D-instanton action *function*. In fact, the integration variables in its matrix model are just matrices rather than fields. However, we will refer to that manifold as vacuum moduli space and to these matrices as fields to keep an analogy with the ref. [9].

³ Hence, when $R^4/\alpha'^2 \ll 1$ and all the curved part of the bulk space-time is confined inside sub-stringy distances. Still, D-instanton can probe it.

Third, it is the limit $N \rightarrow \infty$ with $n \rightarrow \infty$ taken on the "Higgs branch", which is related to the presence of a large number of the D3-branes in the IKKT M-atrrix theory [9, 19]. Thus, even at this point we see that the $N \rightarrow \infty$ D3-branes are equivalent to the $AdS_5 \times S_5$ SUGRA. In our next publication we hope to present a proof of those facts and the statements of the ref. [17] via the methods presented in this note.

2. For the beginning, we explain how a D-instanton can probe an external metric and what is the meaning of the low energy expansion in this context. In early papers on the subject in question, D-branes with the world history were considered as probes. This allowed to use the common way of testing the metric of an external space-time [4]: via the kinetic part in the world-volume theory of a probe. In the D-instanton case, in turn, there is no such a term, because it is just a point in the ambient space. However, in the ref. [12] it was argued that D-branes also can probe an external metric via commutators in their world-volume theory. For example, in the case of $n > 1$ D-instantons one can have the following effective action:

$$S_{inst} = \frac{const \cdot \alpha'^2}{g_s} Tr \left(- G_{\mu\mu'}(A) \cdot G_{\nu\nu'}(A) \cdot [A^\mu, A^\nu] [A^{\mu'}, A^{\nu'}] \right) + \\ + \text{super - partners} + O\left(Tr[.,.]^3\right), \quad A = Tr \sum_{\mu=1}^{10} A_\mu^2. \quad (1)$$

where A_μ , ($\mu = 1, \dots, 10$) are $U(n)$ matrices. Their eigen-values parametrize positions of the D-instantons in the target space with the metric $G_{\mu\nu}$. The first term in this formula is unambiguous.

The contribution $O\left(Tr[A, A]^3\right)$ in eq. (1) includes all higher powers in commutators. They are necessary [12] to maintain an equivalent of the "general covariance"⁴ in this context. This series is an analog of an expansion in powers of derivatives, i.e. of the low energy expansion. In fact, under a T-duality transformation the A_μ matrices are exchanged with differential operators – the covariant derivatives $\partial_\mu - iA_\mu(x)$.

Let us now describe the D-instanton theory we are going to start with. It is the SUSY $U(n)$ matrix model with additional hyper-multiplets describing the N D3-brane background [19]. Concretely, this theory should contain hyper-multiplets both in the adjoint and N in the fundamental representations of the gauge group. Its action can be obtained, via the reduction from four to zero dimensions, in some particular $\mathcal{N} = 2$, $U(n)$ SYM theory. In $\mathcal{N} = 1$ notations the latter has the following Lagrangian ($g_{YM}^2 = g_s$):

$$L = \frac{1}{g_s} Tr Re \left\{ \frac{1}{2} \int d^2\theta W_\alpha W^\alpha + 2 \int d^4\theta H^+ e^{-2A} H + \right. \\ \left. + 2 \sum_{I=1}^2 \int d^4\theta \Phi_I e^{-2A} \Phi_I + \frac{1}{\sqrt{2}} \int d^2\theta H \epsilon_{IJ} [\Phi_I, \Phi_J] + \right. \\ \left. + 2 \sum_{p=1}^N \left(\int d^4\theta Q_p^+ e^{-2A} Q_p + \int d^4\theta \tilde{Q}_p e^{2A} \tilde{Q}_p^+ + \frac{1}{\sqrt{2}} \int d^2\theta \tilde{Q}_p H Q_p \right) \right\}, \quad (2)$$

where A and H are the vector-multiplet (with the super-field strength W_α) and the adjoint hyper-multiplet, respectively. They compose the $\mathcal{N} = 2$ vector-multiplet. Accordingly,

⁴Which is yet to be understood [12] for such a situation. However, it is approximately respected at each order in the expansion (1).

Φ_I are two adjoint $\mathcal{N} = 1$ hyper-multiplets, belonging to the corresponding $\mathcal{N} = 2$ super-field. Besides that, Q_p and $\tilde{Q}_p$ are hyper-multiplets in the fundamental and anti-fundamental representations of the gauge group, respectively. They, in turn, compose the N fundamental $\mathcal{N} = 2$ super-fields.

The theory (2) describes the low energy dynamics on the world-volume of the n D3-branes in the unstable situation – when N D7 branes are present. In fact, the four-dimensional theory (2) has the β -function leading to the Landau pole. Hence, it is not sensible quantum mechanically. However, its reduction to zero dimensions makes a sense and describes a real and stable situation: the D-instantons in the background of the D3-branes.

After the reduction, the D-instanton action in question looks as follows:

$$\begin{aligned}
S = \frac{const \cdot \alpha'^2}{g_s} Tr \left\{ \right. & -\frac{1}{2} [A_a, A_b]^2 - |[A_a, H]|^2 + \lambda \sigma^a [A_a, \bar{\lambda}] + \\
& + \chi \sigma^a [A_a, \bar{\chi}] - i\sqrt{2} ([\lambda, \chi] H^+ + [\bar{\lambda}, \bar{\chi}] H) + \\
& + \frac{1}{2} ([H^+, H] + [\Phi_I^+, \Phi_I] + Q_p^+ \otimes Q_p - \tilde{Q}_p \otimes \tilde{Q}_p^+)^2 - \\
& - 2 \left| \frac{1}{2} \epsilon_{IJ} [\Phi_I, \Phi_J] + \tilde{Q}_p \otimes Q_p \right|^2 - |[A_a, \Phi_I]|^2 + \psi_I \sigma^a [A_a, \bar{\psi}_I] - \\
& - i\sqrt{2} ([\lambda, \psi_I] \Phi_I^+ + [\bar{\lambda}, \bar{\psi}_I] \Phi_I^+) - 2 |[\Phi_I, H]|^2 - \\
& - \sqrt{2} ([\psi_I, \chi] \Phi_I + [\bar{\psi}_I, \bar{\chi}] \Phi_I^+) - \frac{1}{\sqrt{2}} (H \epsilon_{IJ} [\psi_I, \psi_J] + H^+ \epsilon_{IJ} [\bar{\psi}_I, \bar{\psi}_J]) - \\
& - |A_a Q_p|^2 - |A_a \tilde{Q}_p|^2 - 2 |H Q_p|^2 - 2 |H \tilde{Q}_p|^2 + \\
& + \rho_p \sigma^a A_a \bar{\rho}_p + \tilde{\rho}_p \sigma^a A_a \bar{\tilde{\rho}}_p + i\sqrt{2} (Q_p^+ \lambda \rho_p - \bar{\rho}_p \bar{\lambda} Q_p - \tilde{Q}_p^+ \lambda \tilde{\rho}_p + \bar{\tilde{\rho}}_p \bar{\lambda} \tilde{Q}_p) - \\
& - \sqrt{2} (\tilde{\rho}_p \chi Q_p + \tilde{Q}_p \chi \rho_p + \tilde{\rho}_p H \rho_p + \bar{\rho}_p \bar{\chi} \tilde{Q}_p^+ + Q_p^+ \bar{\chi} \tilde{\rho}_p + \bar{\rho}_p H^+ \tilde{\rho}_p) \left. \right\}. \quad (3)
\end{aligned}$$

Here $\sigma^0 = I$ and σ^a , $(a = 1, \dots, 3)$ are the Pauli matrices. While $\otimes$ means the external product over the $U(n)$ color indexes. We have suppressed them along with the spinor ones. Also from now on we assume a summation over all repeated indexes if the inverse is not stated.

In the action (3) we have symbolized bosonic component fields in the same way as the corresponding $\mathcal{N} = 1$ super-fields. Also λ and χ are super-partners of the four real A_a and one complex H scalars. Similarly ψ_I are that of the two complex fields Φ_I . While, ρ_p and $\tilde{\rho}_p$ are super-partners of the N complex scalars Q_p and $\tilde{Q}_p$, respectively. For our purposes, in deriving the action (3), we have assumed that the D3-branes are coinciding and placed at the origin of the six-dimensional transverse space.

Let us explain now the string and geometric meanings of those bosonic fields. The ten hyper-multiplets A_a , $(a = 1, \dots, 4)$, $H = \frac{1}{\sqrt{2}} (A_5 + iA_6)$ and Φ_I , $(I = 1, 2)$ are coming from the strings attached by both their ends to the D-instantons. According to the ref. [20] one can relate the theory (3), taken only for these fields, to the D-instanton action in the flat target space. At the same time the N hyper-multiplets Q_p and $\tilde{Q}_p$ are coming from the strings stretched between the D-instantons and the D3-branes.

Thus, the eigen-values of the six scalars A_m , $(m = 1, \dots, 6)$ parametrize relative

positions of the n D-instantons to the D3-branes. When those fields acquire non-zero "VEVs", the former are separated from the latter. At the same time, the eigen-values of the four scalars Φ_I represent positions of the D-instantons in the four directions tangential to the D3-branes.

To understand the geometric meaning of the Q_p and $\tilde{Q}_p$ fields one can look for the SUSY "vacuum state" of the theory (3). Among the conditions on the state there are as follows:

$$\begin{aligned} [\Phi_I^+, \Phi_I] + Q_p^+ \otimes Q_p - \tilde{Q}_p \otimes \tilde{Q}_p^+ &= 0 \\ \frac{1}{2} \epsilon_{IJ} [\Phi_I, \Phi_J] + \tilde{Q}_p \otimes Q_p &= 0. \end{aligned} \quad (4)$$

They, along with the factorization over the $U(n)$ gauge group, represent the Hyper-Kahler reduction procedure. Which is the main ingredient in the ADHM construction of the n instanton moduli space in the $U(N)$ gauge theory. (See, for example, [21, 9] for the discussion on this point in the context related to our case.) Therefore, while eigen-values of Φ_I correspond to the positions of the n instantons, their sizes and relative color orientations are represented by the fields Q_p and $\tilde{Q}_p$. If the latter acquire non-zero "VEVs", the D-instantons are confined to the D3-brane world-volume. They become ordinary YM instantons [13] with the sizes set by the background of Q_p and $\tilde{Q}_p$.

3. Now we will be looking for the D-instanton effective action on the "Coulomb branch". As was explained above, it is enough for us to consider the terms which are of the quadratic order in commutators.

On the branch of interest we have a non-zero "VEV" for the $\sum_{m=1}^6 A_m^2 \equiv |A|^2$. Via the gauge fixing it can be set to be (no summation over i is assumed in this formula):

$$\left(\sum_{m=1}^6 A_m^2 \right)_{ij} \equiv |A|_{ij}^2 = \frac{r_i^2}{\alpha'^2} \delta_{ij}, \quad (5)$$

where r_i , ($i = 1, \dots, n$) are the distances between the D-instantons and the D3-brane. Eventually, to simplify an interpretation of the final result, we will take all $r_i \rightarrow r$ (see below). At the same time, while $|A|^2$ is diagonal, A_m themselves are assumed to have small non-diagonal parts. Also we consider non-zero background Φ_I fields with the similar conditions on their non-diagonal parts. Thus, we slightly break SUSY and have small but non-zero commutators of the matrices A_m and Φ_I , which is similar to the case of a non-zero velocity of a probe brane as in the ref. [4].

As we see from the eq. (3) the fields Q_p , $\tilde{Q}_p$, ρ_p and $\tilde{\rho}_p$ acquire "masses" (set by r_i) in the background (5). We integrate them out to get the effective action for the "massless" fields A_m and Φ_I . While "massless" fermions λ , χ and ψ_I are set to zero.

For the background in question the action (3) can be rewritten as follows:

$$\begin{aligned} S = S_{cl} + \frac{const \cdot \alpha'^2}{g_s} Tr \Big\{ & FF^+ - \sqrt{2} F \tilde{Q}_p \otimes Q_p - \sqrt{2} F^+ \tilde{Q}_p^+ \otimes Q_p^+ - \\ & - \frac{1}{2} D^2 + D (Q_p^+ \otimes Q_p - \tilde{Q}_p \otimes \tilde{Q}_p^+) - \\ & - [\Phi_I^+, \Phi_I] (Q_p^+ \otimes Q_p - \tilde{Q}_p \otimes \tilde{Q}_p^+) - \epsilon_{IJ} [\Phi_I, \Phi_J] \tilde{Q}_p^+ \otimes Q_p^+ - \end{aligned}$$

$$\begin{aligned}
& -\epsilon_{IJ} [\Phi_I^+, \Phi_J^+] \tilde{Q}_p \otimes Q_p - |A_m Q_p|^2 - |A_m \tilde{Q}_p|^2 + \rho_p \sigma^a A_a \bar{\rho}_p + \tilde{\rho}_p \sigma^a A_a \bar{\tilde{\rho}}_p - \\
& - \sqrt{2} (\tilde{\rho}_p H \rho_p + \bar{\tilde{\rho}}_p H^+ \bar{\rho}_p) \Big\}, \quad (6)
\end{aligned}$$

where S_{cl} is that part of (3) which depends *only* on the A_m , ($m = 1, \dots, 6$) and Φ_I fields and we have explicitly wrote all the terms depending on the fundamental hyper-multiplets. The fields D , F and F^+ in this formula can be integrated out, leading to the *quartic* in Q_p and $\tilde{Q}_p$ contributions from the eq. (3).

After a straightforward Gaussian integration over the fundamental hyper-multiplets and simple manipulations with the logarithms we get the following effective action⁵:

$$\begin{aligned}
S_{eff} \approx & \frac{const \cdot \alpha'^2}{g_s} Tr \left\{ -\frac{1}{2} [A_m, A_n]^2 - |[A_m, \Phi_I]|^2 + \frac{1}{2} [\Phi_I^+, \Phi_I]^2 - \right. \\
& \left. -\frac{1}{2} |\epsilon_{IJ} [\Phi_I, \Phi_J]|^2 + FF^+ - \frac{1}{2} D^2 \right\} - \\
& -\frac{N}{2} Tr \log \left\{ 1 + \frac{\alpha'^4}{r^4} ([\Phi_I^+, \Phi_I] - D)^2 - \frac{\alpha'^4}{r^4} |\epsilon_{IJ} [\Phi_I, \Phi_J] + \sqrt{2} F|^2 + \dots \right\} + \\
& + \frac{N}{2} Tr \log \left\{ 1 - \frac{\alpha'^4}{r^4} [A_m, A_n]^2 + \dots \right\} + \dots, \quad (7)
\end{aligned}$$

where dots stand for higher powers in the commutators and fields F and D . We have explicitly substituted into this formula the S_{cl} , "VEV" of the $|A|^2$ from the eq. (5) and have taken the limit $r_i \rightarrow r$. The first logarithm, in the obtained expression, is the contribution of the bosons Q_p and $\tilde{Q}_p$, while the second one is due to the fermions ρ_p and $\tilde{\rho}_p$.

To get eventually the terms which are quadratic in commutators, one can expand the logarithms from the eq. (7) only to the leading order. In fact, all higher corrections give, after the integration over the F and D fields, contributions to higher terms in the series (1). Thus, performing the expansion and integration over the F and D fields, we get the ultimate answer:

$$\begin{aligned}
S_{eff} \approx & \frac{const \cdot \alpha'^2}{g_s} \left\{ -\frac{1}{2} \left(1 + \frac{R^4}{r^4} \right) Tr [A_m, A_n]^2 - Tr |[A_m, \Phi_I]|^2 + \right. \\
& \left. + \frac{1}{2} \left(1 + \frac{R^4}{r^4} \right)^{-1} Tr \left([\Phi_I^+, \Phi_I]^2 - |\epsilon_{IJ} [\Phi_I, \Phi_J]|^2 \right) \right\} + O(Tr[.,.]^3), \quad (8)
\end{aligned}$$

which is exactly the D-instanton effective action (1) in the BPS 3-brane background:

$$d^2 s = \left(1 + \frac{R^4}{r^4} \right)^{-\frac{1}{2}} \sum_{I=1}^2 d\varphi_I d\bar{\varphi}_I + \left(1 + \frac{R^4}{r^4} \right)^{\frac{1}{2}} \sum_{m=1}^6 da_m da_m, \quad (9)$$

where $r^2 = \sum_{m=1}^6 a_m^2$. Now one can see that when all commutators are equal to zero, i.e. we respect the SUSY, the bosonic and fermionic determinants are perfectly cancel

⁵We do not include the Fadeev-Popov determinant into this expression, because after the gauge fixing (5) it gives no contribution to the terms quadratic in commutators.

between each other. Moreover, because of the general SUSY arguments, the eq. (8) is an exact result at this order of the expansion (1).

4. As we see the D-instantons allow to probe directly an external metric as in the eq. (1) rather than its implicit form, coming from the Born-Infeld action as in [4]. In the latter case one is able to fix an analog of the "Coulomb behavior" ($\sim v^2/r^2$) in the expansion of an effective action in powers of the velocity (v). Which is more or less expectable within the SYM theory. But this raises the question: how one can understand, within this framework, such a global characteristic of the bulk space-time as the horizon? We could hope that simplicity of the D-instanton matrix model, combined with the direct probing of the metric (1), can help in obtaining the horizon directly in the effective action.

5. Author is indebted for valuable discussions to A. Gorsky, A. Losev, A. Morozov, I. Polyubin and especially to A. Gerasimov. This work was partially supported by RFBR 96-021-7230, RFBR 98-02-16575, INTAS 96-538 and RFBR-96-15-96740 for scientific schools.

References

- [1] J. Polchinski, "*TASI Lectures on D-branes*", hep-th/9611050;
J. Polchinski, S. Chaudhuri and C. Johnson, "*Notes on D-branes*", hep-th/9602052.
- [2] M. Douglas, D. Kabat, P. Pouliot and S. Shenker, *Nucl. Phys.*, **B485** (1997) 85.
- [3] A. Sen, *Nucl. Phys.*, **B475** (1996) 562; *Nucl. Phys.*, **B489** (1997) 139; *Nucl. Phys.*, **B498** (1997) 135;
T. Banks, M. Douglas and N. Seiberg, *Phys. Lett.*, **B387** (1996) 278;
M. Douglas, D. Lowe and J. Schwarz, *Phys. Lett.*, **B394** (1997) 297.
- [4] M. Douglas, J. Polchinski and A. Strominger, "*Probing Five-Dimensional Black Holes with D-branes*", hep-th/9703031.
- [5] E. Witten, *Nucl. Phys.*, **B460** (1996) 335.
- [6] M.R. Douglas and G. Moore, "*D-branes, Quivers and ALE Instantons*", hep-th/9603167.
- [7] E. Akhmedov, "*A remark on the AdS/CFT correspondence and the renormalization group flow*", hep-th/9806217, accepted for publication in PLB.
- [8] J.M. Maldacena, "*The Large N Limit of Superconformal Field Theories and Supergravity*", hep-th/9711200.
- [9] E. Witten, *J.High Energy Phys.*, **07** (1997) 003.
- [10] T. Banks, W. Fischler, S. Shenker and L. Susskind, *Phys. Rev.*, **D55** (1997) 5112.
- [11] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiba, *Nucl. Phys.*, **498** (1997) 467.
- [12] M. Douglas, *Nucl. Phys.Proc.Suppl.*, **68** (1998) 381.
- [13] T. Banks and M. Green, *J.High Energy Phys.*, **05** (1998) 002.

- [14] M. Bianchi, M. Green, S. Kovacs and G. Rossi, *J. High Energy Phys.*, **9808** (1998) 013.
- [15] E. Witten, *J. High Energy Phys.*, **9807** (1998) 006.
- [16] I. Kogan and G. Luzon, "*D-instantons on the boundary*", hep-th/9806197.
- [17] M. Dorey, T. Hollowood, V. Khoze, M. Mattis and S. Vandoren, "*Multi-Instantons and Maldacena's Conjecture*", hep-th/9810243.
- [18] C.-S. Chu, P.-M. Ho and Y.-Y. Wu, "*D-Instanton in AdS_5 and Instanton in SYM_4* ", hep-th/9806103.
- [19] M. Douglas and M. Berkooz, *Phys. Lett.*, **B395** (1997) 196.
- [20] H. Itoyama and A. Tokura, *Phys. Rev.*, **D58** (1998) 026002.
- [21] M. Douglas, "*Branes within Branes*", hep-th/9512077.